



FEP Medical Policy Manual

FEP 7.01.158 Balloon Dilation of the Eustachian Tube

Effective Policy Date: January 1, 2021

Related Policies:

Original Policy Date: March 2018

7.01.105 - Balloon Ostial Dilation for Treatment of Chronic and Recurrent Acute Rhinosinusitis

Balloon Dilation of the Eustachian Tube

Description

Description

Eustachian tube dysfunction occurs when the functional valve of the eustachian tube fails to open and/or close properly. This failure is frequently due to inflammation and can cause symptoms such as muffled hearing, ear fullness, tinnitus, and vertigo. Chronic obstructive eustachian tube dysfunction can lead to hearing loss, otitis media, tympanic membrane perforation, and cholesteatomas. Balloon dilation of the eustachian tube is a procedure intended to improve patency by inflating a balloon in the cartilaginous part of the eustachian tube to cause local dilation.

OBJECTIVE

The objective of this evidence review is to determine whether balloon dilation of the eustachian tube improves the net health outcome in patients with chronic obstructive eustachian tube dysfunction.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

POLICY STATEMENT

Balloon dilation of the eustachian tube (BDET) for treatment of chronic obstructive eustachian tube dysfunction may be considered **medically necessary** under the following conditions:

- Adults (age 22 years and older) with symptoms of obstructive eustachian tube dysfunction (aural fullness, aural pressure, otalgia, and/or hearing loss) for 12 months or longer in one or both ears that significantly affects quality of life or functional health status
 - Aural fullness and pressure must be present (see Policy Guidelines)

AND

- The patient has undergone a comprehensive diagnostic assessment; including patient-reported questionnaires, history and physical exam, tympanometry if the tympanic membrane is intact, nasal endoscopy, and comprehensive audiometry, with the following findings:
 - Abnormal tympanogram (Type B or C)
 - Abnormal tympanic membrane (retracted membrane, effusion, perforation, or any other abnormality identified on exam)

AND

- Failure to respond to appropriate medical management of potential co-occurring conditions, if any, such as allergic rhinitis, rhinosinusitis, and laryngopharyngeal reflux, including 4-6 weeks of a nasal steroid spray, if indicated

AND

- Other causes of aural fullness such as temporomandibular joint disorders, extrinsic obstruction of the eustachian tube, superior semicircular canal dehiscence, and endolymphatic hydrops have been ruled out.

AND

- If the patient had a history of tympanostomy tube placement, symptoms of obstructive eustachian tube dysfunction should have improved while tubes were patent

AND

- The patient does not have patulous eustachian tube dysfunction or another contraindication to the procedure (see Policy Guidelines)

AND

- The patient's eustachian tube dysfunction has been shown to be reversible (see Policy Guidelines)

AND

- Symptoms are continuous rather than episodic (e.g., symptoms occur only in response to barochallenge such as pressure changes while flying)

AND

- The patient has not had a previous BDET procedure

Balloon dilation of the eustachian tube is considered **investigational** if the above criteria are not met.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

POLICY GUIDELINES

Symptoms of obstructive eustachian tube dysfunction may include aural fullness, aural pressure, otalgia, and hearing loss. Nearly all patients will have aural fullness and aural pressure. Many patients will have otalgia, but hearing loss may not be present in all patients (e.g., patients with Type C tympanograms).

Contraindications to Balloon Dilation of the Eustachian Tube

- The following patients should not be considered for balloon dilation of the eustachian tube:
 - Patients with patulous eustachian tube dysfunction (ETD)
 - A diagnosis of patulous ETD is suggested by symptoms of autophony of voice, audible respirations, pulsatile tinnitus, and/or aural fullness.
 - Patients with extrinsic reversible or irreversible causes of eustachian tube dysfunction including but not limited to:
 - craniofacial syndromes, including cleft palate spectrum
 - neoplasms causing extrinsic obstruction of the eustachian tube
 - history of radiation therapy to the nasopharynx
 - enlarged adenoid pads
 - nasopharyngeal mass
 - neuromuscular disorders that lead to hypotonia/ineffective eustachian tube dynamic opening
 - systemic mucosal or autoimmune inflammatory disease affecting the mucosa of the nasopharynx and eustachian tube (e.g. Samter"s triad, Wegener"s disease, mucosal pemphigus) that is ongoing/active (i.e. not in remission)
 - Patients with aural fullness but normal exam and tympanogram
 - Patients with chronic and severe atelectatic ears

Reversibility of Eustachian Tube Dysfunction

Reversibility of Eustachian Tube dysfunction can be demonstrated by several means, including any of the following:

- The patient states that they are able to relieve the pressure by performing a Valsalva maneuver to "pop" their ears
- Performing a Valsalva maneuver produces temporary improvement of the patient"s tympanogram to Type A tympanogram
- Performing a Valsalva maneuver causes the member"s middle ear to aerate, which is indicated by the provider visualizing lateral movement of the tympanic membrane on otoscopy

Balloon Dilation of the Eustachian Tube Used in Combination with Other Procedures

- Patients undergoing BDET concurrent with sinus ostial dilation should meet the same diagnostic criteria for BDET as those undergoing BDET alone.
- Patients with a middle ear effusion at the time of BDET may benefit from concurrent myringotomy with or without tympanostomy tube placement

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

FDA REGULATORY STATUS

Table 1. Devices Cleared by the U.S. Food and Drug Administration

Device	Manufacturer	Date Cleared	510(k) No.	Indication
Acclarent Aera Eustachian Tube Balloon Dilation System	Acclarent, Inc.	01/16/2018	K171761	Eustachian tube dilation
Xpress ENT Dilation System	Entellus Medical, Inc.	04/05/2017	K163509	Eustachian tube dilation

In September 2016, the AERA (Acclarent) was granted a de novo 510(k) classification by the U.S. Food and Drug Administration (FDA) (class II, FDA product code: PNZ). The new classification applies to this device and substantially equivalent devices of this generic type. The AERA is cleared for dilating the ET in patients ages 22 and older with persistent eustachian tube dysfunction (ETD).

In December 2016, the XprESS™ ENT Dilation System (Entellus Medical, Plymouth, MN) was cleared for marketing by the FDA through the 510(k) process (K163509). The FDA determined this device was substantially equivalent to existing devices for use in eustachian tube dysfunction (ETD). The predicate devices are XprESS™ Multi-Sinus Dilation System (K152434) and AERA Eustachian Tube Balloon Dilation System.

RATIONALE

Summary of Evidence

For individuals who have chronic obstructive eustachian tube dysfunction despite medical management who receive balloon dilation of the eustachian tube, the evidence includes randomized controlled trials (RCTs), prospective observational studies, case series, and systematic reviews of these studies. Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. Two 6-week randomized controlled trials found more improvement with balloon dilation plus medical management than medical management alone on patient-reported symptoms, ability to perform a Valsalva maneuver, proportion of patients with normalized tympanograms, and otoscopy findings. Durability of these effects was demonstrated at 52 weeks in the uncontrolled extension phase of both RCTs. No serious device- or procedure-related adverse events were reported through 52 weeks of follow-up. Multiple observational studies and case series have reported that patients experienced improvement when comparing symptoms before and after balloon dilation. The evidence is sufficient to determine the effects of the technology on the net health outcome.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

American Academy of Otolaryngology-Head and Neck Surgery Foundation

In 2019, the American Academy of Otolaryngology published a clinical consensus statement on balloon dilation of the eustachian tube (BDET).² The target population was defined as adults ages 18 years or older who are candidates for BDET because of obstructive eustachian tube dysfunction (ETD) in 1 or both ears for 3 months or longer that significantly affects quality of life or functional health status. The expert panel concluded:

- BDET is an option for treatment of patients with obstructive ETD.
- The diagnosis of obstructive ETD should not be made without a comprehensive and multifaceted assessment, including otoscopy, audiometry, and nasal endoscopy.
- BDET is contraindicated for patients diagnosed as having a patulous ETD
- Further study will be needed to refine patient selection and outcome assessment.

The authors emphasized the importance of identifying other potentially treatable causes of ETD, including allergic rhinitis, rhinosinusitis, and laryngopharyngeal reflux, and noted that medical management of these disorders is indicated prior to offering BDET. They also noted that potential risks of BDET that are relevant to patient counseling include bleeding, scarring, infection, development of patulous ETD, and/or the need for additional procedures.

National Institute for Health and Care Excellence

In 2019, the National Institute for Health and Care Excellence (NICE) published updated guidance on BDET.¹¹ The guidance was based on a rapid review of the evidence,¹² and stated, "Evidence on the safety and efficacy of balloon dilation for eustachian tube dysfunction is adequate to support the use of this procedure provided that standard arrangements are in place for clinical governance, consent and audit." NICE standard arrangements recommendations mean that there is enough evidence for doctors to consider the procedure as an option.

The guidance also noted:

- The procedure was not effective in all patients, and there was little evidence on the benefit of repeat procedures.
- The procedure is only indicated for chronic eustachian tube dysfunction refractory to medical treatment.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

REFERENCES

- Schilder AG, Bhutta MF, Butler CC, et al. Eustachian tube dysfunction: consensus statement on definition, types, clinical presentation and diagnosis. *Clin Otolaryngol.* Oct 2015; 40(5): 407-11. PMID 26347263
- Tucci DL, McCoul ED, Rosenfeld RM, et al. Clinical Consensus Statement: Balloon Dilation of the Eustachian Tube. *Otolaryngol Head Neck Surg.* Jul 2019; 161(1): 6-17. PMID 31161864
- Norman G, Llewellyn A, Harden M, et al. Systematic review of the limited evidence base for treatments of Eustachian tube dysfunction: a health technology assessment. *Clin Otolaryngol.* Feb 2014; 39(1): 6-21. PMID 24438176
- Poe DS, Hanna BM. Balloon dilation of the cartilaginous portion of the eustachian tube: initial safety and feasibility analysis in a cadaver model. *Am J Otolaryngol.* Mar-Apr 2011; 32(2): 115-23. PMID 20392533
- Schroder S, Lehmann M, Ebmeyer J, et al. Balloon Eustachian tuboplasty: a retrospective cohort study. *Clin Otolaryngol.* Dec 2015; 40(6): 629-38. PMID 25867023
- Froehlich MH, Le PT, Nguyen SA, et al. Eustachian Tube Balloon Dilation: A Systematic Review and Meta-analysis of Treatment Outcomes. *Otolaryngol Head Neck Surg.* Jun 02 2020: 194599820924322. PMID 32482125
- Poe D, Anand V, Dean M, et al. Balloon dilation of the eustachian tube for dilatory dysfunction: A randomized controlled trial. *Laryngoscope.* May 2018; 128(5): 1200-1206. PMID 28940574
- Meyer TA, O'Malley EM, Schlosser RJ, et al. A Randomized Controlled Trial of Balloon Dilation as a Treatment for Persistent Eustachian Tube Dysfunction With 1-Year Follow-Up. *Otol Neurotol.* Aug 2018; 39(7): 894-902. PMID 29912819
- Anand V, Poe D, Dean M, et al. Balloon Dilation of the Eustachian Tube: 12-Month Follow-up of the Randomized Controlled Trial Treatment Group. *Otolaryngol Head Neck Surg.* Apr 2019; 160(4): 687-694. PMID 30620688
- Cutler JL, Meyer TA, Nguyen SA, et al. Long-term Outcomes of Balloon Dilation for Persistent Eustachian Tube Dysfunction. *Otol Neurotol.* Dec 2019; 40(10): 1322-1325. PMID 31385858
- National Institute for Health and Care Excellence. Balloon dilation for chronic eustachian tube dysfunction. *Interventional procedures guidance [IPG665].* December 2019. <https://www.nice.org.uk/guidance/ipg665>. Accessed August 14, 2020.
- National Institute for Health and Care Excellence. *Interventional procedure overview of balloon dilation for chronic eustachian tube dysfunction.* <https://www.nice.org.uk/guidance/ipg665/documents/overview-2> December 2019. Accessed August 14, 2020.

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
March 2018	New Policy	Policy created with literature review through October 16, 2017. Balloon dilation of the Eustachian tube for treatment of patients with chronic Eustachian tube dilatory dysfunction is considered investigational.
June 2019	Replace policy	Policy updated with literature review through January 11, 2019. references 14-15 added. Policy statement unchanged.
December 2020	Replace policy	Policy updated with literature review through July 12, 2020. references added. Policy statement changed: Balloon dilation of the eustachian tube for treatment of patients with chronic obstructive eustachian tube dysfunction may be considered medically necessary under specified conditions.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.