

Federal Employee Program® 750 9th St NW Washington, D.C. 20001 202.942.1000 Fax 202.942.1125

5.21.126

Section:Prescription DrugsSubsection:Antineoplastic AgentsSubject:Herceptin Hylecta

Effective Date:July 1, 2024Original Policy Date:March 22, 2019Page:1 of 4

Last Review Date: June 13, 2024

Herceptin Hylecta

Description

Herceptin Hylecta (trastuzumab and hyaluronidase-oysk)

Background

Herceptin Hylecta contains trastuzumab and hyaluronidase. Trastuzumab has been shown to inhibit the proliferation of human tumor cells that overexpress HER2. In vitro, trastuzumabmediated antibody-dependent cellular cytotoxicity (ADCC) has been shown to be preferentially exerted on HER2 overexpressing cancer cells compared with cancer cells that do not overexpress HER2. Hyaluronidase has been shown to increase the absorption rate of a trastuzumab product into the systemic circulation (1).

Regulatory Status

FDA-approved indication: Herceptin Hylecta is a combination of trastuzumab, a HER2/neu receptor antagonist, and hyaluronidase, an endoglycosidase, indicated in adults for: (1)

1. The treatment of HER2-overexpressing breast cancer

Herceptin Hylecta carries a boxed warning regarding possible risks for cardiomyopathy, pulmonary toxicity, and embryo-fetal toxicity. Herceptin Hylecta use can result in cardiac failure that manifests as congestive heart failure (CHF) or decreased left ventricular ejection fraction (LVEF), with greatest risk when administered concurrently with anthracyclines (1).

Exposure to Herceptin Hylecta during pregnancy can result in oligohydramnios, in some cases complicated by pulmonary hypoplasia and neonatal death (1).

5.21.126

Section:	Prescription Drugs	Effective Date:	July 1, 2024
Subsection:	Antineoplastic Agents	Original Policy Date:	March 22, 2019
Subject:	Herceptin Hylecta	Page:	2 of 4

The safety and effectiveness of Herceptin Hylecta in pediatric patients less than 18 years of age have not been established (1).

Related policies

Enhertu, Kadcyla, Margenza, Nerlynx, Perjeta, Phesgo, Trastuzumab, Tukysa, Tykerb

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Herceptin Hylecta may be considered **medically necessary** if the conditions indicated below are met.

Herceptin Hylecta may be considered **investigational** for all other indications.

Prior-Approval Requirements

Age 18 years of age or older

Diagnosis

Patient must have the following:

HER2-overexpressing breast cancer

AND ALL of the following:

- 1. HER2 protein overexpression or HER2 gene amplification as confirmed by an FDA-approved test
- 2. Prescriber agrees to monitor for cardiac function and pulmonary toxicity
- 3. Females of reproductive potential will be advised to use effective contraception during treatment and for 7 months following the last dose
- 4. Inadequate treatment response, intolerance, or contraindication to **ONE** of the preferred products (Kanjinti, Ogivri, Ontruzant)

Prior – Approval Renewal Requirements

Age 18 years of age or older

5.21.126

Section:	Prescription Drugs	Effective Date:	July 1, 2024
Subsection:	Antineoplastic Agents	Original Policy Date:	March 22, 2019
Subject:	Herceptin Hylecta	Page:	3 of 4

Diagnosis

Patient must have the following:

HER2-overexpressing breast cancer

AND ALL of the following:

- 1. Prescriber agrees to monitor for cardiac function and pulmonary toxicity
- 2. Females of reproductive potential will be advised to use effective contraception during treatment and for 7 months following the last dose

Policy Guidelines

Pre – PA Allowance

None

Prior - Approval Limits

Duration 12 months

Prior – Approval Renewal Limits

Same as above

Rationale

Summary

Herceptin Hylecta contains trastuzumab and hyaluronidase. Trastuzumab has been shown to inhibit the proliferation of human tumor cells that overexpress HER2. In vitro, trastuzumabmediated antibody-dependent cellular cytotoxicity (ADCC) has been shown to be preferentially exerted on HER2 overexpressing cancer cells compared with cancer cells that do not overexpress HER2. Hyaluronidase has been shown to increase the absorption rate of a trastuzumab product into the systemic circulation. The safety and effectiveness of Herceptin Hylecta in pediatric patients less than 18 years of age have not been established (1).

Prior approval is required to ensure the safe, clinically appropriate, and cost-effective use of Herceptin Hylecta while maintaining optimal therapeutic outcomes.

References

5.21.126

Section:	Prescription Drugs	Effective Date:	July 1, 2024
Subsection:	Antineoplastic Agents	Original Policy Date:	March 22, 2019
Subject:	Herceptin Hylecta	Page:	4 of 4

- 1. Herceptin Hylecta [package insert]. South San Francisco, CA: Genentech, Inc.; February 2019.
- 2. NCCN Drugs & Biologics Compendium[®] Trastuzumab and hyaluronidase-oysk 2024. National Comprehensive Cancer Network, Inc. Accessed on April 30, 2024.

Policy History

Date	Action
March 2019	Addition to PA
June 2019	Annual review
December 2019	Annual review. Addition of requirement to trial preferred product
March 2020	Annual review
June 2020	Annual review
September 2020	Annual review
December 2020	Annual review. Revised requirement to trial preferred products to: "Patient
	must have tried at least two of the preferred products (Herzuma, Kanjinti,
	Ogivri, Ontruzant, Trazimera) unless the patient has a valid medical
	exception"
June 2021	Annual review and reference update
June 2022	Annual review and reference update
June 2023	Annual review and reference update
December 2023	Annual review and reference update. Per FEP, changed preferred
	products to Kanjinti, Ogivri, and Ontruzant. Also removed Medex
	requirements. Added t/f requirement of ONE preferred agent to initiation
March 2024	Annual review and reference update
June 2024	Annual review and reference update
Keywords	

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on June 13, 2024 and is effective on July 1, 2024.