

Federal Employee Program. Blue Cross Blue Shield Association 750 9th St NW, Suite 900 Washington, D.C. 20001 1-800-624-5060

Fax 1-877-378-4727

5.30.036

Section: **Prescription Drugs Effective Date:** October 1, 2024

Subsection: Endocrine and Metabolic Drugs Original Policy Date: February 17, 2017

Subject: Parathyroid Hormone Analogs 1 of 9 Page:

Last Review Date: September 6, 2024

Parathyroid Hormone Analogs

Description

Bonsity* (teriparatide), Forteo (teriparatide), Teriparatide (teriparatide) Tymlos (abaloparatide)

Background

Bonsity (teriparatide), Forteo (teriparatide) and Teriparatide (teriparatide) are synthetic forms of human parathyroid hormone (PTH), which is the primary regulator of bone and mineral metabolism. The pharmacologic activity of teriparatide, which is similar to the physiologic activity of PTH, includes stimulating osteoblast function, increasing gastrointestinal calcium absorption, and increasing renal tubular reabsorption of calcium. Treatment with teriparatide results in increased bone mineral density, bone mass, and strength. In postmenopausal females, teriparatide has been shown to decrease osteoporosis-related fractures (1-3).

Teriparatide (teriparatide) manufactured by Alvogen is not considered a true generic of Forteo. It is a follow-on teriparatide product approved under the 505 (b) (2) regulatory pathway, with Forteo as the reference drug (3).

Tymlos (abaloparatide) is an analog of human parathyroid hormone related protein (PTHrP[1-34]), which acts as an agonist at the PTH1 receptor (PTH1R). This results in stimulation of osteoblast function and increased bone mass (4).

Regulatory Status

FDA-approved indications:

^{*}This medication is included in this policy but is not available on the market as of yet.

5.30.036

Section: Prescription Drugs Effective Date: October 1, 2024

Subsection: Endocrine and Metabolic Drugs Original Policy Date: February 17, 2017

Subject: Parathyroid Hormone Analogs Page: 2 of 9

Bonsity, **Forteo**, and **Teriparatide** are recombinant human parathyroid hormone analogs (1-34), [rhPTH(1-34)] indicated for: (1-3)

1. Treatment of postmenopausal women with osteoporosis at high risk for fracture

- 2. Increase of bone mass in men with primary or hypogonadal osteoporosis at high risk for fracture
- 3. Treatment of men and women with osteoporosis associated with sustained systemic glucocorticoid therapy at high risk for fracture

Tymlos is a human parathyroid hormone related peptide [PTHrP(1-34)] analog indicated for: (4)

- 1. Treatment of postmenopausal women with osteoporosis at high risk for fracture or patients who have failed or are intolerant to other available osteoporosis therapy
- 2. Treatment to increase bone density in men with osteoporosis at high risk for fracture or patients who have failed or are intolerant to other available osteoporosis therapy

Bonsity, Forteo, Teriparatide, and Tymlos no longer carry a boxed warning about the risk of osteosarcoma, however it is still listed as a warning and precaution. Bonsity, Forteo, Teriparatide, and Tymlos should not be prescribed for patients who are at increased baseline risk for osteosarcoma including those with Paget's disease of bone or unexplained elevations of alkaline phosphatase, pediatric and young adult patients with open epiphyses, or prior external beam or implant radiation therapy involving the skeleton (1-4).

Because of the unknown relevance of the rodent osteosarcoma findings to humans, cumulative use of Tymlos for more than 24 months during a patient's lifetime is not recommended. Bonsity, Forteo, and Teriparatide dosing is no longer limited to 24 months of lifetime use. Use of Bonsity, Forteo, or Teriparatide for more than 24 months during a patient's lifetime should only be considered if a patient remains at or has returned to having a high risk for fracture (1-4).

Caution should be used in prescribing Bonsity, Forteo, or Teriparatide in patients with severe renal impairment. In 5 patients with severe renal impairment (CrCl <30 mL/min), the AUC and T1/2 of teriparatide were increased by 73% and 77%, respectively (1-3).

The safety and effectiveness of Bonsity, Forteo, Teriparatide, and Tymlos in pediatric patients has not been established (1-4).

Related policies

Evenity, Prolia, Xgeva

Subsection: Endocrine and Metabolic Drugs Original Policy Date: February 17, 2017

Subject: Parathyroid Hormone Analogs Page: 3 of 9

Policy

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Bonsity, **Forteo**, **Teriparatide**, and **Tymlos** may be considered **medically necessary** if the conditions indicated below are met.

Bonsity, **Forteo**, **Teriparatide**, and **Tymlos** may be considered **investigational** for all other indications.

Prior-Approval Requirements

Age 18 years of age or older

Diagnoses

Tymlos ONLY

Patient must have **ONE** of the following:

- 1. Postmenopausal women with osteoporosis
- 2. Men with osteoporosis

AND ONE of the following:

- a. History of an osteoporotic low trauma fracture of the spine, hip, proximal humerus, pelvis, or distal forearm
- b. Inadequate treatment response, intolerance, or contraindication to oral or injectable bisphosphonate

AND NONE of the following:

- a. Risk for osteosarcoma
- b. Paget's disease
- c. Unexplained elevations of alkaline phosphatase
- d. Prior bone radiation
- e. Bone metastases or a history of skeletal malignancies
- f. Metabolic bone diseases other than osteoporosis
- g. High levels of calcium
- h. Cumulative lifetime therapy with parathyroid hormone analogs exceeds 24 months (see Appendix 1)

Subsection: Endocrine and Metabolic Drugs Original Policy Date: February 17, 2017

Subject: Parathyroid Hormone Analogs Page: 4 of 9

i. Concurrent therapy with other human parathyroid hormone related peptide analogs (see Appendix 1)

 j. Concurrent therapy with another Prior Authorization (PA) medication for osteoporosis (see Appendix 2)

Bonsity, Forteo, and Teriparatide ONLY

Patient must have **ONE** of the following:

1. Postmenopausal women with osteoporosis

AND ONE of the following:

- a. History of an osteoporotic low trauma fracture of the spine, hip, proximal humerus, pelvis, or distal forearm
- b. Inadequate treatment response, intolerance, or contraindication to oral or injectable bisphosphonate
- 2. Primary or hypogonadal osteoporosis in men

AND ONE of the following:

- a. History of an osteoporotic low trauma fracture of the spine, hip, proximal humerus, pelvis, or distal forearm
- b. Inadequate treatment response, intolerance, or contraindication to oral or injectable bisphosphonate
- 3. Osteoporosis associated with sustained systemic glucocorticoid therapy **AND ONE** of the following:
 - a. History of an osteoporotic low trauma fracture of the spine, hip, proximal humerus, pelvis, or distal forearm
 - b. Inadequate treatment response, intolerance, or contraindication to oral or injectable bisphosphonate **AND** the following:
 - i. Currently receiving or will be initiating glucocorticoid therapy

AND ONE of the following for all indications:

- a. Cumulative lifetime therapy with parathyroid hormone analogs does not exceed 24 months (see Appendix 1)
- b. Patient remains at or has returned to having high risk for fracture despite a total of 24 months of use of parathyroid hormones

AND NONE of the following for all indications:

a. Risk for osteosarcoma

Subsection: Endocrine and Metabolic Drugs Original Policy Date: February 17, 2017

Subject: Parathyroid Hormone Analogs Page: 5 of 9

b. Paget's disease

- c. Unexplained elevations of alkaline phosphatase
- d. Prior bone radiation
- e. Bone metastases or a history of skeletal malignancies
- f. Metabolic bone diseases other than osteoporosis
- g. High levels of calcium
- h. Concurrent therapy with other human parathyroid hormone related peptide analogs (see Appendix 1)
- i. Concurrent therapy with another Prior Authorization (PA) medication for osteoporosis (see Appendix 2)

Prior – Approval Renewal Requirements

Age 18 years of age or older

Diagnoses

Tymlos ONLY

Patient must have **ONE** of the following:

- 1. Postmenopausal women with osteoporosis
- 2. Men with osteoporosis

AND NONE of the following:

- a. Risk for osteosarcoma
- b. Paget's disease
- c. Unexplained elevations of alkaline phosphatase
- d. Prior bone radiation
- e. Bone metastases or a history of skeletal malignancies
- f. Metabolic bone diseases other than osteoporosis
- g. High levels of calcium
- h. Cumulative lifetime therapy with parathyroid hormone analogs exceeds 24 months (see Appendix 1)
- i. Concurrent therapy with other human parathyroid hormone related peptide analogs (see Appendix 1)
- j. Concurrent therapy with another Prior Authorization (PA) medication for osteoporosis (see Appendix 2)

5.30.036

Section: Prescription Drugs Effective Date: October 1, 2024

Subsection: Endocrine and Metabolic Drugs Original Policy Date: February 17, 2017

Subject: Parathyroid Hormone Analogs Page: 6 of 9

Bonsity, Forteo, and Teriparatide ONLY

Patient must have **ONE** of the following:

- 1. Postmenopausal women with osteoporosis
- 2. Primary or hypogonadal osteoporosis in men
- 3. Osteoporosis associated with sustained systemic glucocorticoid therapy

AND ONE of the following for all indications:

- a. Cumulative lifetime therapy with parathyroid hormones does not exceed 24 months
- b. Patient remains at or has returned to having high risk for fracture despite a total of 24 months of use of parathyroid hormones

AND NONE of the following:

- a. Risk for osteosarcoma
- b. Paget's disease
- c. Unexplained elevations of alkaline phosphatase
- d. Prior bone radiation
- e. Bone metastases or a history of skeletal malignancies
- f. Metabolic bone diseases other than osteoporosis
- g. High levels of calcium
- h. Concurrent therapy with other human parathyroid hormone related peptide analogs (see Appendix 1)
- i. Concurrent therapy with another Prior Authorization (PA) medication for osteoporosis (see Appendix 2)

Policy Guidelines

Pre - PA Allowance

None

Prior - Approval Limits

<u>Tymlos</u>

Quantity 3 multi-dose prefilled pens per 90 days

Duration 12 months

Bonsity, Forteo, and Teriparatide

Quantity 3 multi-dose prefilled pens per 84 days

Subsection: Endocrine and Metabolic Drugs Original Policy Date: February 17, 2017

Subject: Parathyroid Hormone Analogs Page: 7 of 9

Duration 12 months

Prior - Approval Renewal Limits

Tymlos

Quantity 3 multi-dose prefilled pens per 90 days

Duration 12 months (**Only ONE renewal**)

Bonsity, Forteo, and Teriparatide

Quantity 3 multi-dose prefilled pens per 84 days

Duration 12 months

Rationale

Summary

Bonsity (teriparatide), Forteo (teriparatide), and Teriparatide (teriparatide) are used in the treatment of postmenopausal women with osteoporosis, primary or hypogonadal osteoporosis in men and osteoporosis associated with sustained systemic glucocorticoid therapy. Tymlos (abaloparatide) is used in the treatment of postmenopausal women or in men with osteoporosis. These agents should not be prescribed for patients who are at increased baseline risk for osteosarcoma including those with Paget's disease of bone or unexplained elevations of alkaline phosphatase, pediatric and young adult patients with open epiphyses, or prior external beam or implant radiation therapy involving the skeleton. The safety and effectiveness of Bonsity, Forteo, Teriparatide, and Tymlos in pediatric patients have not been established (1-4).

Prior authorization is required to ensure the safe, clinically appropriate, and cost-effective use of Bonsity, Forteo, Teriparatide, and Tymlos while maintaining optimal therapeutic outcomes.

References

- 1. Bonsity [package insert]. Morristown, NJ: Alvogen, Inc.; November 2023.
- 2. Forteo [package insert]. Indianapolis, IN: Eli Lilly and Company; April 2021.
- 3. Teriparatide [package insert]. Morristown, NJ: Alvogen Inc.; November 2023.
- 4. Tymlos [package insert]. Waltham, MA: Radius Health, Inc.; December 2023.

Policy History

Date Action Reason

February 2017 Addition to PA

Subsection: Endocrine and Metabolic Drugs Original Policy Date: February 17, 2017

Subject: Parathyroid Hormone Analogs Page: 8 of 9

May 2017 Change in policy name from Forteo To Parathyroid Hormone Analogs

Addition of Tymlos (abaloparatide) to policy and no dual therapy with other

human parathyroid hormone related peptide analogs

June 2017 Annual review
September 2017 Annual review
December 2017 Annual review

November 2018 Annual review and reference update

April 2019 Addition of requirement of no concurrent therapy with another PA

osteoporosis medication and addition of Appendices 1 and 2

June 2019 Annual review
November 2019 Addition of Bonsity
December 2019 Annual review

August 2020 Addition of Teriparatide (biosimilar)
September 2020 Annual review and reference update

January 2021 Forteo boxed warning for osteosarcoma removed. Treatment for Forteo

can now extend beyond 24 months if the patient remains at high risk for

fracture or returns to having high risk for fracture

March 2021 Annual review

September 2021 Annual review and reference update

October 2021 Rearranged and reworded criteria requirements to clarify that cumulative

use of parathyroid hormone analogs, other than Forteo, should not exceed

24 months. Revised background, regulatory and summary sections

December 2021 Annual review

January 2022 Regulatory section updated with the removal of Tymlos boxed warning for

osteosarcoma

March 2022 Annual review and reference update

December 2022 Annual review. Changed policy number to 5.30.036

January 2023 Per PI update, added Tymlos indication of men with osteoporosis

March 2023 Annual review

December 2023 Per PI update, Bonsity and Teriparatide boxed warning for osteosarcoma

removed. Treatment for Bonsity and Teriparatide can now extend beyond 24 months if the patient remains at high risk for fracture or returns to

having high risk for fracture

March 2024 Annual review

September 2024 Annual review and reference update

Keywords

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on September 6, 2024 and is effective on October 1, 2024.

5.30.036

Section: Prescription Drugs Effective Date: October 1, 2024

Subsection: Endocrine and Metabolic Drugs Original Policy Date: February 17, 2017

Subject: Parathyroid Hormone Analogs **Page:** 9 of 9

Appendix 1 - List of human parathyroid hormone related peptide analogs

Generic Name	Brand Name
abaloparatide	Tymlos
teriparatide	Bonsity
teriparatide	Forteo
teriparatide	Teriparatide

Appendix 2 - List of PA Osteoporosis Medications

Generic Name	Brand Name
abaloparatide	Tymlos
denosumab	Prolia
romosuzumab-aqqg	Evenity
teriparatide	Bonsity
teriparatide	Forteo
teriparatide	Teriparatide