

Federal Employee Program. Blue Cross Blue Shield Association 750 9th St NW, Suite 900 Washington, D.C. 20001 1-800-624-5060 Fax 1-877-378-4727

# 5.21.017

Section	Prescription Drugs	Effective Date:	April 1, 2025
Subsection:	Antineoplastic Agents	Original Policy Date:	February 28, 2012
Subject:	Erwinaze	Page:	1 of 4

Last Review Date: March 7, 2025

### Erwinaze

Description

Erwinaze (asparaginase Erwinia chrysanthemi)

#### Background

Erwinaze (asparaginase *Erwinia chrysanthemi*) is an asparagine-specific enzyme. L-asparaginase is a tetrameric enzyme consisting of four identical subunits, each having a molecular weight of about 35 kDa. Asparaginase *Erwinia chrysanthemi* catalyzes the deamidation of asparagine to aspartic acid and ammonia, resulting in a reduction in circulating levels of asparagine. The mechanism of action of Erwinaze is thought to be based on the inability of leukemic cells to synthesize asparagine due to lack of asparagine synthetase activity, resulting in cytotoxicity specific for leukemic cells that depend on an exogenous source of the amino acid asparagine for their protein metabolism and survival (1).

Erwinaze (asparaginase *Erwinia chrysanthemi*) is indicated for the treatment of patients with acute lymphoblastic leukemia (ALL) who have developed hypersensitivity to *E. coli*-derived asparaginase as a component of a multi-agent chemotherapeutic regimen (1).

#### **Regulatory Status**

FDA-approved indication: Erwinaze is indicated as a component of a multi-agent chemotherapeutic regimen for the treatment of patients with acute lymphoblastic leukemia (ALL) who have developed hypersensitivity to *E. coli*-derived asparaginase (1).

Erwinaze is contraindicated in patients with a history of pancreatitis, thrombosis, hemorrhagic events, or anaphylaxis reaction with prior L-asparaginase therapy. Discontinue Erwinaze in the event of serious hypersensitivity reactions, including anaphylaxis, and severe or hemorrhagic

# 5.21.017

Section:	Prescription Drugs	Effective Date:	April 1, 2025
Subsection:	Antineoplastic Agents	Original Policy Date:	February 28, 2012
Subject:	Erwinaze	Page:	2 of 4

pancreatitis. Glucose intolerance can occur and, in some cases, may be irreversible. Perform appropriate monitoring and treat hyperglycemia with insulin, as necessary. If thrombosis or hemorrhage occurs discontinue Erwinaze until resolved. Use in pregnant women only if clearly needed. Do not use in lactating women (1).

Related policies		
Asparlas, Oncaspar, Rylaze		
Policy		

This policy statement applies to clinical review performed for pre-service (Prior Approval, Precertification, Advanced Benefit Determination, etc.) and/or post-service claims.

Erwinaze may be considered **medically necessary** if the conditions indicated below are met.

Erwinaze may be considered investigational for all other indications.

### **Prior-Approval Requirements**

#### Diagnosis

Patient must have the following:

- 1. Acute lymphoblastic leukemia (ALL)
  - a. Hypersensitivity to *E. coli*-derived asparaginase
  - b. Prescriber agrees to monitor bilirubin, liver function tests (LFTs), and glucose

#### Prior – Approval Renewal Requirements

Same as above

**Policy Guidelines** 

Pre - PA Allowance None

Prior - Approval Limits Duration 12 months

### 5.21.017

Section:	Prescription Drugs	Effective Date:	April 1, 2025
Subsection:	Antineoplastic Agents	Original Policy Date:	February 28, 2012
Subject:	Erwinaze	Page:	3 of 4

### Prior – Approval Renewal Limits

Same as above

#### Rationale

#### Summary

Erwinaze (asparaginase *Erwinia chrysanthemi*) can be used as a component of a multi-agent chemotherapeutic regimen for the treatment of pediatric and adult patients with acute lymphoblastic leukemia (ALL) who have developed hypersensitivity to native or pegylated *Escherichia coli (E. coli)*-derived asparaginase. Erwinaze is contraindicated in patients with a history of pancreatitis, thrombosis, hemorrhagic events or anaphylaxis reaction with prior L-asparaginase therapy. Erwinaze therapy should be discontinued if the above conditions occur during therapy. Glucose intolerance could occur and may be irreversible (1).

Prior approval is required to ensure the safe, clinically appropriate, and cost-effective use of Erwinaze while maintaining optimal therapeutic outcomes.

#### References

- 1. Erwinaze [package insert]. Palo Alto, CA: Jazz Pharmaceuticals, Inc.; December 2019.
- 2. NCCN Drugs & Biologics Compendium® Asparaginase Erwinia chrysanthemi 2025. National Comprehensive Cancer Network, Inc. Accessed on January 27, 2025.

Policy History	
Date	Action
February 2012 March 2013 March 2014	New Policy Annual editorial and reference update Annual review
March 2015 June 2016	Annual editorial review and reference update Annual editorial review and reference update Policy number changed from 5.04.17 to 5.21.17
June 2017 June 2018 June 2019	Annual review Annual editorial review Annual review
September 2019	Annual review. Added requirement to monitor bilirubin, LFTs, and glucose per FEP

# 5.21.017

Section:	Prescription Drugs	Effective Date:	April 1, 2025
Subsection:	Antineoplastic Agents	Original Policy Date:	February 28, 2012
Subject:	Erwinaze	Page:	4 of 4

June 2020	Annual review and reference update
March 2021	Annual review
September 2021	Annual review and reference update
March 2022	Annual review and reference update
March 2023	Annual review and reference update. Changed policy number to 5.21.017
March 2024	Annual review and reference update
March 2025	Annual review and reference update
Keywords	

This policy was approved by the FEP® Pharmacy and Medical Policy Committee on March 7, 2025 and is effective on April 1, 2025.